


**Mission:** As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally-owned public lands and natural and cultural resources. This includes fostering wise use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also promotes the goals of the Take Pride in America campaign by encouraging stewardship and citizen responsibility for the public lands and promoting citizen participation in their care. The Department also has a major responsibility for American Indian reservation communities and for people who live in the Island Territories under U.S. Administration.



# DUCKS AT A DISTANCE

A WATERFOWL IDENTIFICATION  
GUIDE




### Flyway Councils

In 1952, Flyway Councils were formed in each of the four flyways. The Council in each flyway is made up of representatives from the wildlife agencies of the States in that flyway—one representative from each State. The Councils study flyway problems, develop waterfowl management recommendations, and generally work closely with the U.S. Fish and Wildlife Service in implementing waterfowl management and research programs.



## Administrative Waterfowl Flyways



### Waterfowl Flyways


The term "flyway" has long been used to designate the migration routes of birds. For management purposes, four waterfowl flyways—Pacific, Central, Mississippi, and Atlantic—were established in the United States in 1948. To varying degrees the waterfowl populations using each of these flyways differ in abundance, species composition, migration pathways, and breeding ground origin. There are differences, also, in levels of shooting pressure and harvest.

For the most part flyway boundaries follow State lines. However, the boundary between the Pacific and the Central flyway general follows the Continental Divide.

There are some problems in matching waterfowl migration corridors with flyway boundaries because some species nest and winter in areas that do not occur along a north-south axis. These species cross flyway boundaries during migration. On balance, the present arrangement is useful in that it permits reasonable management of waterfowl. At some future time, it is possible that further rearrangement of boundaries may permit better management of the waterfowl resource.

## Ducks at a Distance

By Bob Hines  
DEPARTMENT OF THE INTERIOR  
U.S. Fish and Wildlife Service




## Identification is Important

Identifying waterfowl gives many hours of enjoyment to millions of people. This guide will help you recognize birds on the wing—it emphasizes their fall and winter plumage patterns as well as size, shape, and flight characteristics. It does not include local names.

Recognizing the species of ducks and geese can be rewarding to birdwatchers and hunters—and the ducks.

Hunters can contribute to their own sport by not firing at those species that are either protected or scarce, and needed as breeders to restore the flocks. It can add to their daily limit; when extra birds of certain species can be taken legally, hunters who know their ducks on the wing come out ahead.

Knowing a mallard from a merganser has another side: gourmets prefer a corn-fed mallard to the fish duck.





## Wetlands Attract Wildlife

There's more than just ducks in our marshes. Knowing and identifying other birds and animals add to the enjoyment of being in a blind.

The same sources of food and shelter that draw waterfowl to ponds and marshes also attract other forms of wildlife.

Protected species are sometimes more numerous than ducks or geese.

Money from Duck Stamp sales is used exclusively to purchase wetlands, preserving areas for ducks, geese, and all wildlife for the enjoyment and pleasure of hunters and non-hunters alike.



## What to Look For


Differences in size, shape, plumage patterns and colors, wing beat, flocking behavior, voice, and habitat—all help to distinguish one species from another.

Flock maneuvers in the air are clues. Mallards, pintails, and wigeon form loose groups; teal and shovelers flash by in small, compact bunches; at a distance, canvasbacks shift from waving lines to temporary V's.

Closer up, individual silhouettes are important. Variations of head shapes and sizes, lengths of wings and tails, and fat bodies or slim can be seen.

Within shotgun range, color areas can be important. Light conditions might make them look different, but their size and location are positive keys. The sound of their wings can help as much as their calls. Flying goldeneyes make a whistling sound; wood ducks move with a swish; canvasbacks make a steady rushing sound. Not all ducks quack; many whistle, squeal, or grunt.

Although not a hard and fast rule, different species tend to use different types of habitat. Puddle ducks like shallow marshes and creeks while divers prefer larger, deeper, and more open waters.



# Eclipse Plumage



Drake: Spring Plumage

Hen



Drake: Full Eclipse



Drakes Emerging from Eclipse



Most ducks shed their body feathers twice each year. Nearly all drakes lose their bright plumage after mating, and for a few weeks resemble females. This hen-like appearance is called the eclipse plumage. The return to breeding coloration varies in species and individuals of each species. Blue-winged teal and shovellers may retain the eclipse plumage until well into the winter.


Wing feathers are shed only once a year; wing colors are always the same.

Drake: Fall Plumage



## COMPARATIVE SIZES OF WATERFOWL

All birds on these pages are drawn to the same scale.



## Puddle Ducks

Puddle ducks are typically birds of fresh, shallow marshes and rivers rather than of large lakes and bays. They are good divers, but usually feed by dabbling or tipping rather than submerging.

The speculum, or colored wing patch, is generally iridescent and bright, and often a telltale field mark.


Any duck feeding in croplands will likely be a puddle duck, for most of this group are sure-footed and can walk and run well on land. Their diet is mostly vegetable, and grain-fed mallards or pintails or acorn-fattened wood ducks are highly regarded as food.







The mallard is our most common duck, found in all flyways. The males are often called "greenheads." The main wintering area is the lower Mississippi basin, and along the gulf coast, but many stay as far north as open waters permits.



**Mallard**  
Length—24"  
Weight—2¾ lbs.

Eclipse Drake

Hen

## White-Fronted Geese

Length—29 in.  
Weight—6¼ lbs.




Adult

Immature

Migrates chiefly in the Central and Pacific flyways but also present in the Mississippi. Rare in the Atlantic Flyway. Appears brownish gray at a distance. Often called "specklebelly". Most distinctive characteristic of the V-shaped flocks is the high pitched call *kow-kow-kow*.

Two races of snow geese are recognized: greater snows along the Atlantic Coast, and lesser snows elsewhere on the continent. Blue geese are a color phase of the lesser snow.



Adult


Immature

Length—29-31 in.  
Weight—6½-7½ lbs.

## Snow Geese



## Canada Geese



Numerous and popular, Canada geese are often called "honkers." Includes several races varying in weight from 3 to over 12 pounds.

All have black heads and necks, white cheeks, similar habitats and voices. Sexes are identical.

## Brant

Length—24–25 in.

Weight—3 ¼–3 ¾ lbs.


Black Brant



Brant

These are sea geese, the blacks wintering south to Baja, California, in the Pacific.

The Atlantic race winters from Virginia northward. Flight is swift, in irregular and changing flock patterns.




Drake



Drake

Hen



Drake

Hen

Flocks often feed in early morning and late afternoon in nearby harvested fields, returning to marshes and creeks to spend the night.


The flight is not particularly rapid. Hens have a loud *quack*; the drone's voice is a low-pitched *kwek-kwek*.

Typical Flock Pattern






These ducks use all four flyways, but are most plentiful in the west. They are extremely graceful and fast fliers, fond of zig-zagging from great heights before leveling off to land. The long neck and tail make them appear longer than mallards, but in body size and weight they are smaller.



**Pintail**  
Length—26"  
Weight—1¾ lbs.

Once thought to be rare, trumpeter swans are slowly increasing in Alaska and on western refuges and parks. Tundra swans are common and increasing. They winter near Chesapeake Bay, San Francisco Bay, Puget Sound and Salton Sea. Occasionally found in fields. Both species are large with pure white plumage.




**Swans**  
Trumpeter—Length—59 in.  
Weight—28 lbs.  
Tundra —Length—52 in.  
Weight—16 lbs.

## Long-tailed Duck

Length—20 ½ in.  
Weight—2 lbs.

Winter Drake



Summer Drake

Winter Hen

A slim, brightly plumaged sea duck. Smaller than the scoters or eiders.


Flight is swift and low with constantly changing flock formations. Ranges along both coasts and the Great Lakes.

One of the most vocal of ducks; drakes have a loud pleasant *caloo, caloo*, constantly heard.

## Harlequin

Length—17 in.  
Weight—1 ½ lbs.

Drake




Eclipse Drake

Hen

Glossy slate-blue plumage enlivened by white stripes and spots give the adult male harlequin a striking appearance. The female resembles a small female scoter. At a distance, both sexes look black. Flight is swift, with abrupt turns. Flocks are small and compact. Ranges both coasts, north from New Jersey and San Francisco. Uncommon.




Drake



Drake

Hen

They are agile on land and often feed in grain fields. The drakes whistle; the hens have a coarse *quack*.



Drake

Hen

Typical Flock Pattern







Gadwalls are most numerous in the Central Flyway, but not too common anywhere. They are often called "gray mallards" or "gray ducks." They are one of the earliest migrants, seldom facing cold weather. They are the only puddle ducks with a white speculum.



**Gadwall**  
Length—21"  
Weight—2 lbs.

Eclipse Drake

Hen




**Common Eider**

Length—23 1/2 in.  
Weight—5 lbs.

Eclipse Drake

Hen

In flight, drakes appear all black except for the flash of the slight gray underwing and the bright yellow swelling at the base of the upper bill. Scoters feed on mollusks, crabs, and some fish and very little vegetation. They are locally known as "coots."



**Black Scoter**

Length—19 1/2 in.  
Weight—2 1/2 lbs.

Immature

Hen

Thick-necked stocky birds, alternately flapping and sailing in flight; flocks string out in a line, close to the water. Occurs in the United States chiefly along New England coasts and occasionally south to New Jersey. Other eiders—king, spectacled and Stellar's—occur in Alaska and are not pictured in this guide. King eiders occasionally are found in north Atlantic coastal waters.

## White-Winged Scoter

Length—21 ½ in.  
Weight—3 ½ lbs.



The three scoters on these two pages are sea ducks, wintering on open coastal waters. White-wings are among the heaviest and largest of all ducks.

## Surf Scoter

Length—19 ½ in.  
Weight—2 lbs.



Like all scoters, these birds move along our coasts in loose flocks, stringing into irregular, wavy lines. Drakes can be distinguished from other scoters by two white patches on their head and the bright color of the bill.

Flight is strong, direct, usually close to the waves.



Small, compact flocks fly swiftly, usually in a direct line. Wingbeats are rapid.

Drakes whistle and *kack-kack*; hens *quack* like a mallard, but softer.




Typical Flock Pattern



## Wigeon

Length—21"  
Weight—1¾ lbs.



These are nervous birds, quick to take alarm. Their flight is fast, irregular, with many twists and turns. In a bunched flock, their movements have been compared to those of pigeons.

When open water is handy, wigeons often raft up offshore until late afternoon when they move to marshes and ponds to feed.



## Whistling Ducks

Length—18-19 in.  
Weight—1¾ lbs.



The trailing legs and rounded wings of these slow flying ducks makes them look bigger than they are.

Both species are primarily Mexican. In the U.S., the black-bellied is found only in south Texas and Louisiana. The fulvous also occurs there and in Florida with occasional stragglers further north along both coasts and the Mississippi Valley. The fulvous is the more common of the two species in the United States.

Sexes are alike. Both species have shrill whistling calls.





# Hooded Merganser

Length—18 in.  
Weight—1 ½ lbs.

Eclipse Drake



Drake



Hen



Drake



Hen



Often seen in pairs, or very small flocks. Short rapid wingstrokes create an impression of great speed.


Winters in the inland waters of all coastal States; seldom goes to salt water.

Voice: Seldom heard in fall.



Hen

Drake



Drake



Drake



Hen



Drake



Hen


The white belly and forewing are very showy in the air. Drakes whistle; hens have a loud *kaow* and a lower *qua-awk*.

Typical Flock Pattern



## Shoveler

Length—19 1/2"  
Weight—1 1/2 lbs.



Eclipse Drake

Hen

Shovelers, 'spoonbills' to many, are early migrants, moving out at the first frost. The largest numbers are in the Central and Pacific flyways.

The usual flight is steady and direct. When startled, the small flocks twist and turn in the air like teal.



Hen

Drake

## Common Merganser

Length—25 1/2 in.  
Weight—2 1/2 lbs.



Drake

Hen

This species is larger than the red-breasted merganser, and is one of the largest of our ducks. It is one of the last to migrate south, and is more common than the red-breasted merganser on inland waters.

Flocks move in "follow the leader" style, low over the water.

The only call seems to be a startled croak.

Typical Flock Pattern

Hen



Drake



Hen



Drake




Hen

Drake

## Red-Breasted Merganser

Length—23 in.  
Weight—2 ½ lbs.

Eclipse Drake



Hen

Drake

Drake



Hen



Drake



Hen



Typical Flock Pattern



Hen

Drake

These birds winter most abundantly in coastal waters, including the Gulf of Mexico, and to a lesser extent, the Great Lakes.

Their flight, strong and direct, is usually low over the water. They are difficult to distinguish in flight from the common merganser.

Voice: Seldom heard.



Drake



Drake



Hen

They are not highly regarded as table birds, because one third of the usual diet is animal matter.



Drake



Hen

Drakes call *woh-woh* and *took-took*; the hen's *quack* is feeble.

Typical Flock Pattern





# Blue-Winged Teal

Length—16"  
Weight—15 oz.




Eclipse Drake

Hen

Their small size and twisting turning flight gives the illusion of great speed. The small, compact flocks commonly fly low over the marshes, and often take the hunter by surprise. They are more vocal than most ducks—their high-pitched peeping and nasal quacking is commonly heard in spring and to a lesser extent in fall. These teal are among the first ducks to migrate each fall, and one of the last in the spring.

Drake



Hen



Drake



Hen



Hen

Drake



Summer Drake

Drake



Hen



Drake



Hen



They are early to mid-fall migrants. Drakes often cock their tails upright at an angle, the only species to habitually do so. Both hens and drakes are silent in the fall.

Typical Flock Pattern



## Ruddy

Length—15 ½ in.  
Weight—1 ⅓ lbs.



The ruddy duck often dives or swims away from danger rather than flying. When flying, their small wings stroke so fast they resemble bumblebees.



Sexes Similar



## Cinnamon Teal





Blue-Winged Drake

In the Pacific Flyway, cinnamon teal are far more common than blue-wings. The hens look alike and the habits of both species are similar.

The pale blue forewing patch is the best field mark, as drakes are usually in eclipse until January or longer.

Drakes have a whistling *peep*; hens utter a low *quack*.

Typical Flock Pattern



# Green-Winged Teal

Length—15 in.  
Weight—14 oz.

Eclipse Drake



Hen

Quite hardy—some birds stay as far north as open water is found.  
The smallest and one of the most common of our ducks. Their tiny size gives the impression of great speed, but mallards can fly faster. Their flight is often low, erratic, with the entire flock twisting and turning as one unit.

Hen

Drake



Drake



Drake



Hen



Largest concentrations are on both seacoasts and along the Gulf of Mexico. Inland, they will remain as far north as open water permits.

Usually silent. Drakes *squeak* and have a guttural note; hens *quack* weakly.

Drake



Hen



Typical Flock Pattern





## Bufflehead

Length—14 ½ in  
Weight—1 lb.



Hen

Eclipse Drake

Stragglers migrate south in mid-fall, but the largest numbers move just ahead of freezeup. Most flocks in feeding areas are small—5 or 6 birds, with more hens and immatures than adult drakes.

Very small size, bold black and white color pattern, and low, swift flight are field marks. Unlike most divers, they can fly straight up from a watery takeoff.



Hen

Drake



Drake



Drake

Hen



Drake



Hen

They nest as far north as Alaska, and migrate in all four flyways. Early fall drakes are usually still in full eclipse plumage.

Drakes whistle and twitter; hens have a slight *quack*.

Typical Flock Pattern



## Wood Duck

Length—18 ½ in.  
Weight—1 ½ lbs.



Found in all flyways; most numerous in the Atlantic and Mississippi flyways and fewest in the Central. They are early migrants; most of them have left the northern States by mid-November.

Frequents wooded streams and ponds; perches in trees. Flies through thick timber with speed and ease and often feeds on acorns, berries, and grapes on the forest floors.



Hen

Drake



Barrow's Drake

Common Drake


Barrow's goldeneye, predominantly a westerner, is less wary than the common goldeneye.

Hens of both species are look-alikes.

Drakes have a piercing *spee-speer*—hens a low *quack*. Both are usually quiet.



Drake



Hen



Typical Flock Pattern



Hen

## Goldeneye

Common—Length—19 in.  
Weight—2¼ lbs.

Barrow's—Length—19 in.  
Weight—2¾ lbs.



Hen Both Species


Common Eclipse Drake

These are active, strong-winged fliers moving singly or in small flocks, often high in the air. Distinctive wing-whistling sound in flight has earned the name of whistlers.

Goldeneyes generally move south late in the season; most of them winter on coastal waters and the Great Lakes. Inland, they like rapids and fast water.

Barrow's

Common



Hen

Drake

Hen


Drake



Drake



Drake



Hen



Drake



Hen

Flight is swift and direct; flocks are usually small.

In the air, their wings make a rustling, swishing sound. Drakes call *hoo-w-ett*, often in flight; hens have a *cr-r-ek* when frightened.

Typical Flock Pattern





# Black Duck

Length—24 in.  
Weight—2¾ lbs.



A bird of the eastern States, primarily the Atlantic Flyway and, to a lesser extent, the Mississippi.

Shy and wary, regarded as the

wariest of all ducks.

Often seen in company of mallards, but along the Atlantic coast frequents the salt marshes and ocean much more than mallards. Flight is swift, usually in small flocks.

White wing lining in contrast to very dark body plumage is a good identification clue.

The hen's *quack* and the drake's *kwek-kwek* are duplicates of the mallards.

Typical Flock Pattern




Similar Sexes



Greater Scaup Drake

Lesser Scaup Drake





Both species migrate late, sometimes just before freezeup.

Flock movements are rapid, often erratic, usually in compact groups.

Hens are silent; drake lesser scaup *purrr*; drake greater scaup have a discordant *scaup, scaup*.


Typical Flock Pattern



## Scaup

Greater—Length—18 ½ in.  
Weight—2 lbs.

Lesser—Length—17 in.  
Weight—1 ⅞




Except for the wing marks, greater and lesser scaup appear nearly identical in the field.

The light band near the trailing edges of the wings runs almost to the tip in the greater scaup, but only about half way in the lesser.

Greater scaup prefer large open water areas; lesser scaup often use marshes and ponds.

Lesser

Greater



## Diving Ducks

Diving ducks frequent the larger, deeper lakes and rivers, and coastal bays and inlets.

The colored wing patches of these birds lack the brilliance of the speculums of puddle ducks. Since many of them have short tails, their huge, paddle feet may be used as rudders in flight, and are often visible on flying birds. When launching into flight, most of this group patter along the water before becoming airborne.

They feed by diving, often to considerable depths. To escape danger, they can travel great distances underwater, emerging only enough to show their head before submerging again.

Their diets of fish, shellfish, mollusks, and aquatic plants make them second choice, as a group, for sportsmen. Canvasbacks and redheads fattened on eel grass or wild celery are notable exceptions.

Since their wings are smaller in proportion to the size and weight of their bodies, they have a more rapid wingbeat than puddle ducks.



# Canvasback

Length—22 in.  
Weight—3 lbs.



Eclipse Drake

Hen

Normally late to start south, canvasbacks migrate in lines and irregular V's. In feeding areas, compact flocks fly in indefinite formations. Their wingbeat is rapid and noisy; their speed is the swiftest of all our ducks.



Hen

Drake



Drake

Fly as small flocks in open formation; often land without circling. Drakes *purr*; hens are usually silent.



Typical Flock Pattern

Drake

Hen

Drake



# Ringneck

Length—17 in.  
Weight—2 ½ lbs.

Eclipse Drake



Hen

Similar in appearance to scaups, but more often found in fresh marshes and wooded ponds. In flight, the dark wings are different from the white-edged wings of scaup.

Faint brown ring on drake's neck never shows in the field; light bands at tip and base of bill are conspicuous.



Hen

Drake



Drake

Feeding behavior is highly variable. In some areas they feed at night and spend the day rafted up in open waters; in other areas they feed inshore mornings and evenings.

On the water, body size and head shape distinguish them from scaups and redheads.

Drakes *croak*, *peep*, and *growl*; hens have a mallard-like *quack*.



Drake

Hen



Drake

Hen




Typical Flock Pattern



# Redheads

Length—20 in.  
Weight—2 1/2 lbs.



Range coast to coast, with the largest numbers in the Central Flyway. Migratory flocks travel in V's; move in irregular formations over feeding areas. Often found associating with canvasbacks. In the air, they give the impression of always being in a hurry.



Usually spend the day in large rafts in deep water; feed morning and evening in shallower sections. Drakes *purr* and *meow*; hens have a loud *squak*, higher than a hen mallard's.



Typical Flock Pattern